
β Tubulin **Rabbit** Polyclonal antibody


目 录 号：ANT327
规　　格：50μl
储　　存：-20℃/一年

组　　成：PBS with 0.02% sodium azide and 50% glycerol, pH7.3
来　　源：Rabbit / IgG
应　　用：WB, IHC, IF/ICC, IF-P, FC (Intra), IP, ELISA
浓　　度：0.6mg/ml
反应种属：human, mouse, rat
条带大小：50KDa
稀　　释：WB: 1:2000-1:12000
IP: 0.5-4.0 ug for 1.0-3.0mg of total protein lysate
IHC: 1:50-1:500
IF-P: 1:50-1:500
IF/ICC: 1:200-1:800
FC (Intra): 0.25 ug per 10^6 cells in a 100μl suspension
为获得最佳实验效果，请根据具体实验调整抗体最佳工作浓度。


相关产品：
ANT321 β-actin Antibody (Mab)
ANT322 β-actin Antibody (Pab)
ANT324 GAPDH Antibody (Mab)
ANT325 GAPDH Antibody (Pab)
ANT326 β-tubulin (Mab)
ANT328 alpha-tubulin (Mab)
ANT329 alpha-tubulin (Pab)




Various lysates were subjected to SDS PAGE followed by western blot with 10094-1-AP (Beta Tubulin antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 10094-1-AP (beta Tubulin antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.




IP result of anti-Beta Tubulin (IP:10094-1-AP, 3ug; Detection:10094-1-AP 1:1000) with mouse brain tissue lysate 7500ug.




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 10094-1-AP (Beta Tubulin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 10094-1-AP (Beta Tubulin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using Beta Tubulin antibody (10094-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Beta Tubulin antibody (10094-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug Beta Tubulin Polyclonal antibody (10094-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug rabbit IgG isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.